


Protein Naked Cuticle Homolog 1 (NKD1) Antibody

Catalogue No.: abx027501



In the mouse, Nkd is a Dishevelled (see DVL1; MIM 601365) binding protein that functions as a negative regulator of the Wnt (see WNT1; MIM 164820) beta-catenin (see MIM 116806) Tcf (see MIM 602272) signaling pathway.

Target:	Protein Naked Cuticle Homolog 1 (NKD1)
Clonality:	Polyclonal
Reactivity:	Mouse
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	WB: 1/1000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	KLH-conjugated synthetic peptide between 83-111 amino acids from the N-terminal region of human NKD1.
Isotype:	IgG
Form:	Liquid
Purification:	Purified through a protein A column, followed by peptide affinity purification.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q969G9 (UniProt , ExPASy)
KEGG:	hsa:85407

Datasheet

Version: 3.0.0

Revision date: 21 Jun 2025



String: [9606.ENSF00000268459](#)

Molecular Weight: Calculated MW: 52.3 kDa

Buffer: PBS containing 0.09% sodium azide.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only