



Tropomyosin Beta Chain (TPM2) Antibody


Catalogue No.: abx002226




Western blot analysis of various lysates using TPM2 Antibody at 1/400 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.




Immunohistochemistry analysis of paraffin-embedded Rat heart using TPM2 Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human stomach using TPM2 Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse heart using TPM2 Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.



Immunoprecipitation analysis of 300 µg extracts of Mouse skeletal muscle cells using 3 µg TPM2 antibody. Western blot was performed from the immunoprecipitate using TPM2 antibody at a dilution of 1/1000.

Datasheet

Version: 5.0.0
Revision date: 30 Jul 2025



TPM2 Antibody is a Rabbit Polyclonal antibody against TPM2. This gene encodes beta-tropomyosin, a member of the actin filament binding protein family, and mainly expressed in slow, type 1 muscle fibers. Mutations in this gene can alter the expression of other sarcomeric tropomyosin proteins, and cause cap disease, nemaline myopathy and distal arthrogryposis syndromes. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Target:	Tropomyosin Beta Chain (TPM2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IHC, IF/ICC, IP
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/1000, IHC-P: 1/50 - 1/100, IF/ICC: 1/50 - 1/100, IP: 0.5 µg - 4 µg antibody per 400 µg - 600 µg extracts of whole cells. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Synthetic peptide corresponding to TPM2. The exact sequence is proprietary.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P07951 (UniProt , ExPASy)
Gene Symbol:	TPM2
GeneID:	7169
NCBI Accession:	NP_003280.2
KEGG:	hsa:7169
Molecular Weight:	Calculated MW: 33 kDa Observed MW: 36 kDa
Buffer:	PBS, pH 7.3, containing 0.01% thimerosal, 50% glycerol.

Datasheet

Version: 5.0.0

Revision date: 30 Jul 2025



Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only