



Nuclear Factor 1 C-Type (NFIC) Antibody

Catalogue No.: abx028560



NFIC recognizes and binds the palindromic sequence 5'-TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.

Target: Nuclear Factor 1 C-Type (NFIC)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1/1000, IF/ICC: 1/10 - 1/50. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic peptide between 180-208 amino acids from the Central region of human NFIC.

Isotype: IgG

Form: Liquid

Datasheet

Version: 4.0.0
Revision date: 01 Jun 2025



Purification:	Purified through a protein A column, followed by peptide affinity purification.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P08651 (UniProt , ExPASy)
NCBI Accession:	NP_001231931.1, NP_001231933.1, NP_001231934.1, NP_005588.2, NP_995315.1
KEGG:	hsa:4782
String:	9606.ENSP00000396843
Molecular Weight:	Calculated MW: 55.7 kDa
Buffer:	PBS containing 0.09% sodium azide.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only