


General Transcription Factor IIIC Subunit 2 (GTF3C2) Antibody

Catalogue No.: abx028690



GTF3C2 is required for RNA polymerase III-mediated transcription. Component of TFIIC that initiates transcription complex assembly on tRNA and is required for transcription of 5S rRNA and other stable nuclear and cytoplasmic RNAs. May play a direct role in stabilizing interactions of TFIIC2 with TFIIC1.

Target:	General Transcription Factor IIIC Subunit 2 (GTF3C2)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	WB: 1/1000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	KLH-conjugated synthetic peptide between 141-170 amino acids from the N-terminal region of human GTF3C2.
Isotype:	IgG
Form:	Liquid
Purification:	Purified through a protein A column, followed by peptide affinity purification.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q8WUA4 (UniProt , ExPASy)
Gene Symbol:	GTF3C2

Datasheet

Version: 2.0.0

Revision date: 26 Jun 2025



KEGG: hsa:2976

String: [9606.ENSP00000352536](#)

Molecular Weight: Calculated MW: 101 kDa

Buffer: PBS containing 0.09% sodium azide.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only