

Interferon-Stimulated Gene 20 kDa Protein (ISG20) Antibody

Catalogue No.: abx028929



Exonuclease with specificity for single-stranded RNA and, to a lesser extent for DNA. Degrades RNA at a rate that is approximately 35-fold higher than its rate for single-stranded DNA. Involved in the antiviral function of IFN against RNA viruses.

| | |
|-------------------------------|--|
| Target: | Interferon-Stimulated Gene 20 kDa Protein (ISG20) |
| Clonality: | Polyclonal |
| Reactivity: | Human |
| Tested Applications: | ELISA, WB |
| Host: | Rabbit |
| Recommended dilutions: | WB: 1/1000. Optimal dilutions/concentrations should be determined by the end user. |
| Conjugation: | Unconjugated |
| Immunogen: | KLH-conjugated synthetic peptide between 29-58 amino acids from the Central region of human ISG20. |
| Isotype: | IgG |
| Form: | Liquid |
| Purification: | Purified through a protein A column, followed by peptide affinity purification. |
| Storage: | Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles. |
| UniProt Primary AC: | Q96AZ6 (UniProt , ExPASy) |
| KEGG: | hsa:3669 |

Datasheet

Version: 5.0.0

Revision date: 28 Sep 2025



String: [9606.ENSF00000306565](#)

Molecular Weight: Calculated MW: 20.4 kDa

Buffer: PBS containing 0.09% sodium azide.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only