



Serine-Threonine Kinase Receptor-Associated Protein (STRAP) Antibody

Catalogue No.:abx031805





17 (-) (+)



The SMN complex plays an essential role in spliceosomal snRNP assembly in the cytoplasm and is required for pre-mRNA splicing in the nucleus. STRAP may play a role in the cellular distribution of the SMN complex.

Target: Serine-Threonine Kinase Receptor-Associated Protein (STRAP)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, IHC, FCM

Host: Rabbit

Recommended dilutions: WB: 1/1000, IHC-P: 1/50 - 1/100, FCM: 1/10 - 1/50. Not tested in IHC-F. Optimal

dilutions/concentrations should be determined by the end user.

Datasheet

Version: 3.0.0 Revision date: 03 May 2025



Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic peptide between 242-269 amino acids from the C-terminal region of

human STRAP.

Isotype: IgG

Form: Liquid

Purification: Purified Rabbit Polyclonal Antibody.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q9Y3F4 (UniProt, ExPASy)

String: <u>9606.ENSP00000392270</u>

Molecular Weight: Calculated MW: 38.4 kDa

Buffer: PBS containing 0.09% sodium azide.

Specificity: Predicted to react with Mouse, Rat, Cow and Chicken STRAP.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.