


Bcl2-Associated Agonist of Cell Death Phospho-Tyr110 (BAD pY110) Antibody

Catalogue No.: abx032127



The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq].

Target: Bcl2-Associated Agonist of Cell Death Phospho-Tyr110 (BAD pY110)

Clonality: Polyclonal

Target Modification: Tyr110

Modification: Phosphorylation

Reactivity: Human

Tested Applications: ELISA, DB

Host: Rabbit

Recommended dilutions: DB: 1/500. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y110 of human BAD.

Isotype: IgG

Form: Liquid

Datasheet

Version: 2.0.0
Revision date: 23 Oct 2025



Purification:	Purified through a protein A column, followed by two-step phosphospecific peptide affinity purification.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q92934 (UniProt , ExPASy)
KEGG:	hsa:572
String:	9606.ENSP00000378040
Molecular Weight:	Calculated MW: 18.4 kDa
Buffer:	PBS containing 0.09% sodium azide.
Specificity:	Predicted to react with Mouse and Rat BAD.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only