


RNA Polymerase-Associated Protein CTR9 Homolog (CTR9) Antibody

Catalogue No.: abx002638



Western blot analysis of various lysates using CTR9 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 30s.

CTR9 Antibody is a Rabbit Polyclonal antibody against CTR9. CTR9, parafibromin (CDC73; MIM 607393), LEO1 (MIM 610507), and PAF1 (MIM 610506) form the PAF protein complex, which associates with the RNA polymerase II subunit POLR2A (MIM 180660) and with a histone methyltransferase complex (Rozenblatt-Rosen et al., 2005).

Target:	RNA Polymerase-Associated Protein CTR9 Homolog (CTR9)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/1000 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1024-1173 of human CTR9.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q6PD62 (UniProt , ExPASy)
Gene Symbol:	CTR9

Datasheet

Version: 4.0.0

Revision date: 05 Mar 2025



GeneID: [9646](#)

NCBI Accession: NP_055448.1

KEGG: hsa:9646

String: [9606.ENSP00000355013](#)

Molecular Weight: Calculated MW: 134 kDa
Observed MW: 145 kDa

Buffer: PBS, pH 7.3, containing 0.01% thimerosal, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only