



Upstream Stimulatory Factor 2 (USF2) Antibody

Catalogue No.: abx034010



USF2 is a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. This protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E-box motifs.

Target:	Upstream Stimulatory Factor 2 (USF2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse
Tested Applications:	ELISA, WB, IHC
Host:	Rabbit
Recommended dilutions:	WB: 1/1000, IHC-P: 1/50 - 1/100. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	KLH-conjugated synthetic peptide between 218-246 amino acids from the Central region of human USF2.
Isotype:	IgG
Form:	Liquid

Datasheet

Version: 4.0.0

Revision date: 06 Sep 2025



Purification:	Purified through a protein A column, followed by peptide affinity purification.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q15853 (UniProt , ExPASy)
KEGG:	hsa:7392
String:	9606.ENSP00000222305
Molecular Weight:	Calculated MW: 37 kDa
Buffer:	PBS containing 0.09% sodium azide.
Specificity:	Predicted to react with Rat USF2.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only