



Signal Transducer And Activator of Transcription 3 (STAT3) Antibody


Catalogue No.: abx132219




Western blot analysis of recombinant Human STAT3.



IHC-P analysis of Human kidney Tissue, with DAB staining.



IHC-P analysis of human stomach tissue, with DAB staining.



IHC-P analysis of human liver tissue, with DAB staining.

Signal Transducer And Activator Of Transcription 3 (STAT3) Antibody is a Rabbit Polyclonal against Signal Transducer And Activator Of Transcription 3 (STAT3).

Target: Signal Transducer And Activator of Transcription 3 (STAT3)

Clonality: Polyclonal

Datasheet

Version: 3.0.0
Revision date: 15 Aug 2025



Reactivity:	Human, Mouse, Rat, Cow
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant Signal Transducer And Activator Of Transcription 3 (STAT3) (Arg335-Trp546) expressed in E. coli.
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P40763 (UniProt , ExPASy)
KEGG:	hsa:6774
String:	9606.ENSP00000264657
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.