


Stromal Cell Derived Factor 4 (SDF4) Antibody (Biotin)

Catalogue No.: abx433478



Biotinylated abx431576 (0.1 µg/ml) staining of Jurkat lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.

SDF4 Antibody is a goat polyclonal antibody against SDF4.

Target: Stromal Cell Derived Factor 4 (SDF4)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: P-ELISA, WB

Host: Goat

Recommended dilutions: P-ELISA: 1/64000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Biotin

Immunogen: Internal region, 161-175 AA: C-EYKVKFLASKGHSEK

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: SDF4

GeneID: [51150](#) [20318](#) [155173](#)

NCBI Accession: NP_057631.1, NP_057260.2

Datasheet

Version: 3.0.0

Revision date: 21 May 2025



Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only