

Activated Leukocyte Cell Adhesion Molecule / CD166 (ALCAM) Antibody (FITC)

Catalogue No.: abx273869



Fluorescence emission spectra of FITC.

Activated Leukocyte Cell Adhesion Molecule (ALCAM) Antibody (FITC) is a Rabbit Polyclonal antibody conjugated to FITC against Activated Leukocyte Cell Adhesion Molecule (ALCAM).

Target:	Activated Leukocyte Cell Adhesion Molecule / CD166 (ALCAM)
Research Area:	Cellular Differentiation and Adhesion, Tumor Immunity
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.5-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	FITC
Excitation/Emission:	499/515
Laser Line:	488
Isotype:	IgG
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography.
Storage:	Aliquot and store at 4°C. Avoid exposure to light. Avoid repeated freeze/thaw cycles.

Datasheet

Version: 2.0.0

Revision date: 19 Nov 2025



Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only