



Thymic Stromal Lymphopoietin (TSLP) Antibody


Catalogue No.: abx174774



Immunohistochemistry analysis of Human Liver Tissue using Thymic Stromal Lymphopoietin Antibody (30 µg/ml) and HRP-conjugated Goat Anti-Mouse antibody ([abx400001](#), 2 µg/ml), with DAB staining.



Immunohistochemistry analysis of Human Spleen Tissue using Thymic Stromal Lymphopoietin Antibody (30 µg/ml) and HRP-conjugated Goat Anti-Mouse antibody ([abx400001](#), 2 µg/ml), with DAB staining.



Immunohistochemistry analysis of Human Kidney Tissue using Thymic Stromal Lymphopoietin Antibody (30 µg/ml) and HRP-conjugated Goat Anti-Mouse antibody ([abx400001](#), 2 µg/ml), with DAB staining.

Thymic Stromal Lymphopoietin (TSLP) Antibody is a Mouse Monoclonal antibody against Thymic Stromal Lymphopoietin (TSLP).

Target: Thymic Stromal Lymphopoietin (TSLP)

Clonality: Monoclonal

Clone: C523

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Mouse

Datasheet

Version: 3.0.0
Revision date: 12 Sep 2025



Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-30 µg/ml, IF/ICC: 5-30 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx168695](#) - Recombinant TSLP (Tyr29-Gln159) expressed in E. coli

Isotype: IgG₁ Kappa

Form: Liquid

Purification: Purified by Protein A and Protein G affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Buffer: 0.01 M PBS, pH 7.4, containing 0.02% NaN₃, 50% glycerol.

Specificity: Cross-reacts with Pig TSLP.

Concentration: 0.61 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only