



Wntless Wnt Ligand Secretion Mediator (WLS) Antibody


Catalogue No.: abx338537



WB analysis of HepG2 whole cell lysate, using WLS antibody (5.4 µg/ml). Predicted band size: 63, 52 kDa. Observed band size: 52 kDa.



IHC-P analysis of human kidney tissue. Antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0).



IHC-P analysis of human adrenal gland tissue. Antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0).

WLS Antibody is a Rabbit Polyclonal against WLS.

Target: Wntless Wnt Ligand Secretion Mediator (WLS)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/5000, IHC: 1/200 - 1/500. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 1.0.0
Revision date: 24 May 2025



Immunogen:	Recombinant Human Protein wtless homolog protein (123-232 AA).
Isotype:	IgG
Form:	Liquid
Purity:	> 95%
Purification:	Purified by Protein G.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q5T9L3 (UniProt , ExPASy)
Gene Symbol:	WLS
GeneID:	79971
KEGG:	hsa:79971
String:	9606.ENSP00000346829
Buffer:	0.01 M PBS, pH 7.4, 0.03% Proclin-300 and 50% Glycerol.
Concentration:	2.73 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.