



Somatoliberin (GHRH) Antibody

Catalogue No.:abx004085



Western blot analysis of various lysates using GHRH Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 180s.



Immunofluorescence analysis of NTERA-2 cells using GHRH Antibody at dilution of 1/100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.

GHRH Antibody is a Rabbit Polyclonal antibody against GHRH. The protein encoded by this gene belongs to the glucagon family and is a preproprotein that is produced in the hypothalamus. The preproprotein is cleaved to form a 44 aa factor, also called somatocrinin, that acts to stimulate growth hormone release from the pituitary. Variant receptors for somatocrinin have been found in several types of tumors, and antagonists of these receptors can inhibit the growth of the tumors. Defects in this gene are a cause of dwarfism, while hypersecretion of the encoded protein is a cause of gigantism.

Target: Somatoliberin (GHRH)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

Recommended dilutions: ELISA: 1 μg/ml, WB: 1/500 - 1/1000, IF/ICC: 1/50 - 1/200. Optimal dilutions/concentrations should

be determined by the end user.

Conjugation: Unconjugated

Immunogen: Synthetic peptide corresponding to GHRH. The exact sequence is proprietary.

Isotype: IgG

Datasheet

Version: 3.0.0 Revision date: 21 Oct 2025



Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P01286 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: GHRH

GeneID: <u>2691</u>

NCBI Accession: NP_066567.1

KEGG: hsa:2691

String: <u>9606.ENSP00000362716</u>

Molecular Weight: Calculated MW: 12 kDa

Observed MW: 12 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.