


Human Bruton'S Tyrosine Kinase (Btk) Protein

Catalogue No.:abx065610



SDS-PAGE analysis of Human Btk Protein.

Recombinant Bruton'S Tyrosine Kinase (Btk) is a recombinant Human protein produced in a Prokaryotic expression system (E. coli).

| Target: | Bruton'S Tyrosine Kinase (Btk) | |
|-----------------------------------|---|--|
| Origin: | Human | |
| Expression: | Recombinant | |
| Tested Applications: WB, SDS-PAGE | | |
| Host: | E. coli | |
| Conjugation: | Unconjugated | |
| Form: | Lyophilized | |
| Purity: | > 95% | |
| Reconstitution: | To keep the original salt concentration, we recommend reconstituting to the original concentration prior to lyophilization (see Concentration) in ddH_2O . If a lower concentration is required, dilute in PBS, pH 7.4. If a higher concentration is required, the product can be reconstituted directly in PBS, pH 7.4, though please note that this will change the overall salt concentration. The stock concentration should be between 0.1-1.0 mg/ml. Do not vortex. | |
| Storage: | Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw cycles. | |
| UniProt Primary AC: | Q06187 (<u>UniProt</u> , <u>ExPASy</u>) | |
| KEGG: | hsa:695 | |
| String: | <u>9606.ENSP00000483570</u> | |
| | | |

Abbexa BV, Leiden, NL Website: www.abbexa.com · Email: info@abbexa.com Datasheet Version: 3.0.0 Revision date: 14 Apr 2025



| Molecular Weight: | Calculated MW: 57.9 kDa |
|--------------------|--------------------------------|
| | Observed MW (SDS-PAGE): 55 kDa |
| Sequence Fragment: | Glu396-Ser659 |

 Sequence:
 EIDPK DLTFLKELGT GQFGVVKYGK WRGQYDVAIK MIKEGSMSED EFIEEAKVMM NLSHEKLVQL

 YGVCTKQRPI FIITEYMANG CLLNYLREMR HRFQTQQLLE MCKDVCEAME YLESKQFLHR

 DLAARNCLVN DQGVVKVSDF GLSRYVLDDE YTSSVGSKFP VRWSPPEVLM YSKFSSKSDI

 WAFGVLMWEI YSLGKMPYER FTNSETAEHI AQGLRLYRPH LASEKVYTIM YSCWHEKADE

 RPTFKILLSN ILDVMDEES

Tag: N-terminal His tag and GST tag

Buffer: Prior to lyophilization: PBS, pH 7.4, containing 0.01% Sarcosyl, 1 mM DTT, 5% Trehalose and Proclin-300.

Activity: Not tested

Concentration: Prior to lyophilization: 200 µg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.