


Human Calcitonin Gene Related Peptide 2 / CGRP2 (CALCB) Protein

Catalogue No.:abx065658



SDS-PAGE analysis of Human Calcitonin Protein.

Human Calcitonin Gene Related Peptide (CGRP) is a recombinant Human protein produced in a Prokaryotic expression system (E. coli).

Target: Calcitonin Gene Related Peptide 2 / CGRP2 (CALCB)

Origin: Human

Expression: Recombinant

Tested Applications: WB, SDS-PAGE

Host: E. coli

Conjugation: Unconjugated

Form: Lyophilized

Purity: > 95%

Reconstitution: To keep the original salt concentration, we recommend reconstituting to the original concentration prior

to lyophilization (see Concentration) in ddH₂O. If a lower concentration is required, dilute in 20 mM Tris,

150 mM NaCl, pH 8.0. If a higher concentration is required, the product can be reconstituted directly in 20 mM Tris, 150 mM NaCl, pH 8.0, though please note that this will change the overall salt

concentration. The stock concentration should be between 0.1-1.0 mg/ml. Do not vortex.

Storage: Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw

cycles.

UniProt Primary AC: P10092 (UniProt, ExPASy)

KEGG: hsa:797

String: 9606.ENSP00000433490

Datasheet

Version: 3.0.0 Revision date: 26 Jun 2025



Molecular Weight: Calculated MW: 15.0 kDa

Observed MW (SDS-PAGE): 19 kDa

Possible reasons why the actual band size differs from the predicted band size:

- 1. Splice variants. Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge. The composition of amino acids may affect the charge of the protein.
- 3. Post-translational modification. Phosphorylation, glycoslyation, methylation etc. may affect the band size.
- 4. Post-translational cleavage. Many proteins are synthesised as pro-proteins, and then cleaved to give
- 5. Polymerisation of the target protein. Dimerisation, multimerisation etc. will increase the band size observed.

Sequence Fragment: Met1-Ala127

Sequence: MGFRKFSPFL ALSILVLYQA GSLQAAPFRS ALESSPDPAT LSKEDARLL AALVQDYVQM

KASELKQEQE TQGSSSAAQK RACNTATCVT HRLAGLLSRS GGMVKSNFVP TNVGSKAFGR

RRRDLQA

Tag: N-terminal His tag

Buffer: Prior to lyophilization: 20 mM Tris, 150 mM NaCl, pH 8.0, containing 1 mM EDTA, 1 mM DTT, 0.01%

Sarcosyl, 5% Trehalose and Proclin-300.

Activity: Not tested

Concentration: Prior to lyophilization: 200 µg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC

OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.