



Human Desmoglein 1 (DSG1) Protein

Catalogue No.: abx066312



SDS-PAGE analysis of recombinant Human Desmoglein 1 Protein.



Gene sequencing extract of recombinant Human Desmoglein 1 Protein.

Desmoglein 1 (DSG1) is a recombinant Human protein produced in a Prokaryotic expression system (E. coli).

Target: Desmoglein 1 (DSG1)

Origin: Human

Expression: Recombinant

Tested Applications: WB, SDS-PAGE

Host: E. coli

Conjugation: Unconjugated

Form: Lyophilized

Purity: > 90%

Reconstitution: To keep the original salt concentration, we recommend reconstituting to the original concentration prior to lyophilization (see Concentration) in ddH₂O. If a lower concentration is required, dilute in 20 mM Tris, 150 mM NaCl, pH 8.0. If a higher concentration is required, the product can be reconstituted directly in 20 mM Tris, 150 mM NaCl, pH 8.0, though please note that this will change the overall salt concentration. The stock concentration should be between 0.1-1.0 mg/ml. Do not vortex.

Datasheet

Version: 4.0.0
Revision date: 14 May 2025



Storage: Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q02413 ([UniProt](#), [ExPASy](#))

KEGG: hsa:1828

String: [9606.ENSP00000257192](#)

Molecular Weight: Calculated MW: 30.3 kDa
Observed MW (SDS-PAGE): 40 kDa
Possible reasons why the actual band size differs from the predicted band size:

1. Splice variants. Alternative splicing may create different sized proteins from the same gene.
2. Relative charge. The composition of amino acids may affect the charge of the protein.
3. Post-translational modification. Phosphorylation, glycosylation, methylation etc. may affect the band size.
4. Post-translational cleavage. Many proteins are synthesised as pro-proteins, and then cleaved to give the active form.
5. Polymerisation of the target protein. Dimerisation, multimerisation etc. will increase the band size observed.

Sequence Fragment: Glu710-Val961

Sequence: E GRPSNDCLLI YDIEGVGSPA GSVGCCSFIG EDLDDSFLLT LGPKFKKLAD ISLGKESYPD
LDPSWPPQST EPVCLPQETE PVVSGHPPIS PHFGTTTVIS ESTYPSGPGV LHPKPILDPL
GYGNVTVTES YTTSIDLKPS VHVHDNRPAS NVVVTERTVVG PISGADLHGM LEMPDLRDGS
NVIVTERVIA PSSSLPTSLT IHHPRESSNV VVTERVIQPT SGMIGSLSMH PELANAHNVI
VTERVVSGAG V

Tag: N-terminal His tag

Buffer: Prior to lyophilization: 20 mM Tris, 150 mM NaCl, pH 8.0, containing 0.01% Sarcosyl, 5% Trehalose.

Activity: Not tested

Concentration: Prior to lyophilization: 200 µg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.