



Voltage-Dependent Anion Channel 1 (VDAC1) Antibody

Catalogue No.: abx239848



WB analysis of HEK-293 cells, using VDAC1 antibody (1/1000 dilution).



IHC-P analysis of human colon tissue, using VDAC1 antibody (1/50 dilution).

Voltage-Dependent Anion Channel 1 (VDAC1) Antibody is a Rabbit Polyclonal antibody against Voltage-Dependent Anion Channel 1 (VDAC1).

Target: Voltage-Dependent Anion Channel 1 (VDAC1)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/100 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: voltage-dependent anion channel 1

Isotype: IgG

Form: Liquid

Datasheet

Version: 2.0.0
Revision date: 02 Sep 2025



Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P21796 (UniProt , ExPASy)
KEGG:	hsa:7416
String:	9606.ENSP00000265333
Molecular Weight:	Observed MW: 35 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.