


Human Inducible T-Cell Co Stimulator Ligand (ICOSLG) Protein

Catalogue No.: abx067185



SDS-PAGE analysis of Human ICOSLG Protein.

Inducible T-Cell Co Stimulator Ligand (ICOSLG) protein is a Recombinant protein expressed in *E. coli*.

Target: Inducible T-Cell Co Stimulator Ligand (ICOSLG)

Research Area: Signal Transduction, Tumor Immunity, Infection Immunity, Immunology

Origin: Human

Expression: Recombinant

Tested Applications: WB, SDS-PAGE

Host: *E. coli*

Conjugation: Unconjugated

Form: Lyophilized

Activity: Not tested

Purity: > 95%

Reconstitution: To keep the original salt concentration, we recommend reconstituting to the original concentration prior to lyophilization (see Concentration) in ddH₂O. If a lower concentration is required, dilute in PBS, pH 7.4. If a higher concentration is required, the product can be reconstituted directly in PBS, pH 7.4, though please note that this will change the overall salt concentration. The stock concentration should be between 0.1-1.0 mg/ml. Do not vortex.

Storage: Store at 2-8°C for up to one month. For long-term storage, store at -80°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: O75144 ([UniProt](#), [ExPASy](#))

Datasheet

Version: 2.0.0

Revision date: 10 Oct 2025



Molecular Weight: Calculated MW: 29.1 kDa

Sequence Fragment: Asp19-Ser258

Tag: N-terminal His tag

Buffer: Prior to lyophilization: PBS, pH 7.4, containing 0.01% Sarcosyl, 1 mM DTT, 5% Trehalose and Proclin-300.

Endotoxin Level: < 1.0 EU per 1 µg (determined by the LAL method)

Concentration: Prior to lyophilization: 200 µg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only