


TP63 Antibody

Catalogue No.: abx243053



Western Blot

Positive WB detected in: PC3 whole cell lysate


All lanes: TP63 antibody at 2.1 µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 66, 63, 52, 56, 45, 58, 47, 68, 57, 77 KDa

Observed band size: 77 KDa



Immunohistochemistry analysis of TP63 Antibody diluted at 1/210 and staining in paraffin-embedded human pancreatic cancer. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

TP63 Antibody is a Monoclonal Antibody against TP63.

Target:	TP63
Clonality:	Monoclonal
Reactivity:	Human
Expression:	Recombinant
Tested Applications:	ELISA, WB, IHC
Host:	Rabbit

Recommended dilutions: WB: 1/500 - 1/5000, IHC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: A synthesized peptide derived from human TP63.

Isotype: IgG

Form: Liquid

Datasheet

Version: 3.0.0

Revision date: 22 Sep 2025



Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20 °C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9H3D4 (UniProt , ExPASy)
Buffer:	PBS, pH 7.4, 150 mM NaCl, 0.02% sodium azide and 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only