



CCL19 Antibody

Catalogue No.: abx243056



Immunocytochemistry analysis of CCL19 Antibody diluted at 1/155 and staining in paraffin-embedded human tonsil tissue. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry analysis of CCL19 Antibody diluted at 1/155 and staining in paraffin-embedded human appendix tissue. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

CCL19 Antibody is a Monoclonal Antibody against CCL19.

Target:	CCL19
Clonality:	Monoclonal
Clone:	P238
Reactivity:	Human
Expression:	Recombinant
Tested Applications:	ELISA, IHC
Host:	Rabbit

Recommended dilutions: IHC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: A synthesized peptide derived from human CCL19.

Isotype: IgG

Datasheet

Version: 2.0.0

Revision date: 06 Sep 2025



Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20 °C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q99731 (UniProt , ExPASy)
KEGG:	hsa:6363
String:	9606.ENSP00000308815
Buffer:	PBS, pH 7.4, 150 mM NaCl, 0.02% sodium azide and 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only