



C-C Chemokine Receptor Type 9 (CCR9) Antibody


Catalogue No.:abx243073



WB analysis of SH-SY5Y, HepG2, Jurkat, HL-60, HeLa, U87 cell lysates and Mouse brain tissue lysate using CCR9 antibody (1 µg/ml) and a Goat anti-Rabbit IgG antibody (1/50000 dilution). Calculated MW: 41 kDa, 43 kDa, Observed MW: 43 KDa.



IF analysis of HepG2 cells using CCR9 Antibody (1/33 dilution) and a Goat anti-Rabbit IgG H&L antibody conjugated to AF488. DAPI was used as a nuclear stain.



IP analysis of CCR9 in HL-60 lysate. Lane 1: Negative control Rabbit IgG antibody with a Protein G antibody conjugated to HRP (1/2000 dilution), Lane 2: 500 µg HL-60 lysate with CCR9 antibody (3 µg/ml) with a Protein G antibody conjugated to HRP (1/2000 dilution) and Lane 3: 20 µg HL-60 lysate.

C-C Chemokine Receptor Type 9 (CCR9) Antibody is a Monoclonal Antibody against CCR9.

Target:	C-C Chemokine Receptor Type 9 (CCR9)
Clonality:	Monoclonal
Clone:	V147
Reactivity:	Human, Mouse
Expression:	Recombinant
Tested Applications:	ELISA, WB, IF/ICC, IP
Host:	Rabbit

# Datasheet

Version: 3.0.0

Revision date: 06 May 2025



**Recommended dilutions:** WB: 1/500 - 1/5000, IF/ICC: 1/20 - 1/200, IP: 1/200 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

**Conjugation:** Unconjugated

**Immunogen:** A synthesized peptide derived from Human CCR9.

**Isotype:** IgG

**Form:** Liquid

**Purification:** Purified by affinity chromatography.

**Storage:** Aliquot and store at -20 °C. Avoid repeated freeze/thaw cycles.

**UniProt Primary AC:** P51686 ([UniProt](#), [ExPASy](#))

**KEGG:** hsa:10803

**String:** [9606.ENSP00000350256](#)

**Buffer:** PBS, pH 7.4, 150 mM NaCl, 0.02% sodium azide and 50% glycerol.

**Concentration:** 1.5 mg/ml

**Note:** THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.