



Leukemia Inhibitory Factor (LIF) Antibody


Catalogue No.: abx433588



IHC-P analysis of paraffin embedded Human Testis tissue using LIF antibody (6 µg/ml).



Negative Control showing staining of paraffin embedded Human Testis tissue, with no primary antibody.



Flow cytometry analysis of paraformaldehyde fixed HepG2 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

LIF Antibody is a Goat Polyclonal Antibody against LIF.

Target: Leukemia Inhibitory Factor (LIF)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: P-ELISA, IHC, FCM

Host: Goat

Recommended dilutions: P-ELISA: 1/128000, IHC: 6 µg/ml, FCM: 10 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Immunogen: [abx616964](#) - Internal region: RLCSKYHVGHVDT

Datasheet

Version: 1.0.0

Revision date: 20 Mar 2025



Epitope Region:	Internal region.
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	LIF
GeneID:	3976
NCBI Accession:	NP_002300.1
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only