


Potassium Voltage-Gated Channel Subfamily J Member 11 (KCNJ11) Antibody

Catalogue No.: abx004413



Western blot analysis of extracts of various cell lines using KCNJ11 Antibody (1/1000 dilution).

KCNJ11 Antibody is a Rabbit Polyclonal antibody against KCNJ11. Potassium channels are present in most mammalian cells, where they participate in a wide range of physiologic responses. The protein encoded by this gene is an integral membrane protein and inward-rectifier type potassium channel. The encoded protein, which has a greater tendency to allow potassium to flow into a cell rather than out of a cell, is controlled by G-proteins and is found associated with the sulfonylurea receptor SUR. Mutations in this gene are a cause of familial persistent hyperinsulinemic hypoglycemia of infancy (PHHI), an autosomal recessive disorder characterized by unregulated insulin secretion. Defects in this gene may also contribute to autosomal dominant non-insulin-dependent diabetes mellitus type II (NIDDM), transient neonatal diabetes mellitus type 3 (TNDM3), and permanent neonatal diabetes mellitus (PNDM). Multiple alternatively spliced transcript variants that encode different protein isoforms have been described for this gene.

Target: Potassium Voltage-Gated Channel Subfamily J Member 11 (KCNJ11)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: WB

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant fusion protein corresponding to human KCNJ11

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Datasheet

Version: 2.0.0
Revision date: 24 Jun 2025



Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q14654 ([UniProt](#), [ExPASy](#))

Gene Symbol: KCNJ11

GeneID: [3767](#)

NCBI Accession: NP_000516.3

KEGG: hsa:3767

String: [9606.ENSP00000345708](#)

Molecular Weight: Calculated MW: 33 kDa/43 kDa
Observed MW: 31-43 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.