Datasheet

Version: 4.0.0 Revision date: 10 Apr 2025



SARS-CoV-2 S1 Protein RBD (Gamma P.1 Variant)

Catalogue No.:abx160086

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2/COVID-19) Spike S1 Protein Receptor-Binding Domain (RBD) is a recombinant protein expressed in Insect cells.

The SARS-CoV-2 Spike Protein (S protein) is a viral protein that allows the entry of SARS-CoV-2 into human cells. The protein forms trimers on the viral capsid and binds to human Angiotensin Converting Enzyme 2 (ACE2) located on the cell surface. The protein has a cleavage site between the S1 and S2 subunits that is targeted by the human enzyme Furin, and it may also cause the development of a syncytium (cell fusion). Antibodies to S protein can prevent viral entry as well as target the virus for further immune action.

The Gamma (P.1) variant was first identified in Brazil. It contains the K417T, E484K and N501Y mutations in the spike protein's receptor-binding domain.

Target: Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) S1 Protein RBD (Gamma

B.1.1.28.1 Variant)

Target Modification: Lys417Thr, Glu484Lys, Asn501Tyr

Modification: Mutation

Origin: Virus

Expression: Recombinant

Tested Applications: ELISA, WB

Host: Insect

Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Form: Liquid

Purity: $\geq 90\%$ (SDS-PAGE)

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Molecular Weight: Calculated MW: 26 kDa

Observed MW (12% reducing SDS-PAGE): ~28 kDa

Datasheet

Version: 4.0.0 Revision date: 10 Apr 2025



Sequence Fragment: 319-541 AA.

Tag: C-terminal His tag

Buffer: PBS, pH 7.4. Does not contain preservatives.

Concentration: 2.9 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY, NOT FOR USE IN DIAGNOSTIC.

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.

This product is shipped with dry ice.

Directions for use: Centrifuge before opening to ensure complete recovery of vial contents.

