

Chemically Competent Cell

Catalogue No.: abx098066

Chemically Competent Cell is designed for chemical transformation of DNA, with a transformation efficiency of over 10^8 cfu/ μ g DNA (tested by pUC19 plasmid DNA). It is suitable for blue/white selection, routine cloning, and is the lowest homologous recombination favourable for plasmid DNA preparation.

The genotype is: *endA1 recA1 gyrA96 thi-1 hsdR17* (r_k^- , m_k^+) *relA1 supE44 D (lac-proAB)* [*F'**traD36 proAB laqI* Δ M15].

The 1 ml size consists of 10 \times 100 μ l Competent Cells, 20 μ l (0.1 ng/ μ l) Control Plasmid pUC19, and 10 ml SOC Medium.

Target: Chemically Competent Cell

Storage: Store at -70 °C for up to 6 months. Do not store in liquid nitrogen. Avoid repeated freeze/thaw cycles.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.
This product is shipped with dry ice.

Directions for use:

Recommended Protocol:

1. Equilibrate a water bath to 42 °C. Bring a vial of SOC medium or LB medium to room temperature. Warm selective plates to 37 °C for 30 minutes.
2. Thaw 100 μ l of Chemically Competent Cell on ice. Aliquot 50 μ l of cells into a pre-chilled 1.5 ml tube, then add target DNA (1-5 μ l). Do not mix by pipetting up and down. Leave on ice for 30 minutes.
3. Heat-shock the cells for 45 seconds at 42 °C, without shaking. Immediately transfer to the tube to ice. Leave on ice for 2 minutes without shaking.
4. Add 500 μ l of prewarmed SOC medium or LB medium (without antibiotics) into the tube. Mix well and shake at 37 °C for 1 hour at 200 RPM for cell recovery and expression of antibiotic resistance.
5. Spread 20-200 μ l from each transformation vial onto a pre-warmed selective plate. Cells that have not been plated can be stored at 4 °C and plated the next day if required.
6. Invert the plates and incubate at 37 °C overnight.
7. Select colonies and analyse by restriction enzyme digestion, PCR or sequencing.

Note:

Higher efficiency transformation can be achieved by transforming cells immediately following thawing. Avoid repeated thawing. All samples and reagents require gentle handling throughout the entire procedure.