


Datasheet

Version: 3.0.0


Revision date: 02 Jun 2025

Pepsin Antibody


Catalogue No.: abx100019



Western blot analysis of Human Urine.



IHC-P analysis of Human Stomach Tissue, with DAB staining.



IHC-P analysis of Human Prostate Gland Tissue, with DAB staining.

Polyclonal Antibody to Pepsin.

Target: Pepsin

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 3.0.0

Revision date: 02 Jun 2025



Immunogen:	Native Protein Pepsin
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	0.83 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only