



Prorelaxin H2 (RLN2) Antibody


Catalogue No.: abx100410




Western blot analysis of Human Placenta Tissue.



Western blot analysis of recombinant Human RLN2.



IHC-P analysis of Human Liver Tissue, with DAB staining.



IHC-P analysis of Human Liver Cancer Tissue, with DAB staining.

Polyclonal Antibody to Prorelaxin H2 (RLN2).

Target: Prorelaxin H2 (RLN2)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 2.0.0
Revision date: 01 Oct 2025



Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	RLN2 (Met1-Cys186)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P04090 (UniProt , ExPASy)
Gene Symbol:	RLN2
GeneID:	6019
OMIM:	179740
HGNC:	10027
KEGG:	hsa:6019
Ensembl:	ENSG00000107014
String:	9606.ENSP00000371040
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.