



NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2) Antibody


Catalogue No.: abx100644




Western blot analysis of (1) Rat Brain Tissue and (2) Mouse Brain Tissue.




Western blot analysis of recombinant Rat SIRT2.



IHC-P analysis of Rat Testis Tissue, with DAB staining.



IHC-P analysis of Rat Stomach Tissue, with DAB staining.



IHC-P analysis of Rat Intestine Tissue, with DAB staining.

Datasheet

Version: 3.0.0
Revision date: 08 Sep 2025



Polyclonal Antibody to NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2).

Target:	NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2)
Research Area:	Signal Transduction, Developmental Science
Clonality:	Polyclonal
Reactivity:	Rat
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	SIRT2 (Glu28-Leu303)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q5RJQ4 (UniProt , ExPASy)
Gene Symbol:	SIRT2
KEGG:	rno:361532
String:	10116.ENSARNOP00000059450
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.