



NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6) Antibody

Catalogue No.: abx100653



Western blot analysis of recombinant Human SIRT6.



IHC-P analysis of Bowels tissue, with DAB staining.

Polyclonal Antibody to NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6).

Target: NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6)

Research Area: Signal Transduction

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1-5 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: SIRT6 (Met1-Ser355)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Datasheet

Version: 4.0.0
Revision date: 13 Oct 2025



Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q8N6T7 ([UniProt](#), [ExPASy](#))

Gene Symbol: SIRT6

GeneID: [51548](#)

OMIM: [606211](#)

HGNC: 14934

KEGG: hsa:51548

Ensembl: ENSG00000077463

String: [9606.ENSP00000337332](#)

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only