



NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6) Antibody


Catalogue No.: abx100654



WB analysis of recombinant Mouse NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6).



IHC-P analysis with DAB staining, of Mouse brain tissue, using NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6) antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis with DAB staining, of Mouse subcutaneous tumor tissue, using NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6) antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml). PBS was used instead of the primary antibody as a negative control.

Polyclonal Antibody to NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6).

Target: NAD-Dependent Protein Deacetylase Sirtuin-6 (SIRT6)

Research Area: Signal Transduction

Clonality: Polyclonal

Reactivity: Mouse

Tested Applications: WB, IHC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Datasheet

Version: 5.0.0
Revision date: 17 Sep 2025



Conjugation:	Unconjugated
Immunogen:	abx069109 - recombinant Mouse SIRT6 (Met1-Ser334) expressed in E. coli.
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P59941 (UniProt , ExPASy)
Gene Symbol:	SIRT6
GeneID:	50721
KEGG:	mmu:50721
Ensembl:	ENSMUSG00000034748
String:	10090.ENSMUSP00000048971
Buffer:	PBS, pH 7.4, containing 0.02% NaN ₃ , 50% glycerol.
Specificity:	Cross-reacts with Rat SIRT6.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.