



Anamorsin (CIAPIN1) Antibody


Catalogue No.: abx004844



Western blot analysis of various lysates using CIAPIN1 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 10s.



Immunofluorescence analysis of U2OS cells using CIAPIN1 Antibody. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 200 µg extracts of 293T cells, using 3 µg CIAPIN1 antibody. Western blot was performed from the immunoprecipitate using CIAPIN1 antibody at a dilution of 1/1000.

CIAPIN1 Antibody is a Rabbit Polyclonal antibody against CIAPIN1. CIAPIN1 is a cytokine-induced inhibitor of apoptosis with no relation to apoptosis regulatory molecules of the BCL2 (MIM 151430) or CASP (see MIM 147678) families. Expression of CIAPIN1 is dependent on growth factor stimulation (Shibayama et al., 2004).

Target: Anamorsin (CIAPIN1)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IF/ICC, IP

Host: Rabbit

Datasheet

Version: 5.0.0
Revision date: 26 Oct 2025



Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/2000, IF/ICC: 1/50 - 1/200, IP: 0.5 µg - 4 µg antibody per 200 µg - 400 µg extracts of whole cells. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to CIAPIN1. The exact sequence is proprietary.

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q6FI81 ([UniProt](#), [ExPASy](#))

Gene Symbol: CIAPIN1

GeneID: [57019](#)

OMIM: [608943](#)

NCBI Accession: NP_064709.2

HGNC: 28050

KEGG: hsa:57019

Ensembl: ENSG00000005194

String: [9606.ENSP00000377914](#)

Molecular Weight: Calculated MW: 34 kDa
Observed MW: 39 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.