



Oncostatin-M (OSM) Antibody


Catalogue No.: abx100844




Western blot analysis of recombinant Mouse OSM.




IHC-P analysis of Mouse Testis Tissue, with DAB staining.



IHC-P analysis of Mouse Heart Tissue, with DAB staining.



IHC-P analysis of Mouse Stomach Tissue, with DAB staining.



IHC-P analysis of Mouse Liver Tissue, with DAB staining.

Datasheet

Version: 5.0.0
Revision date: 30 Aug 2025



Polyclonal Antibody to Oncostatin-M (OSM).

Target:	Oncostatin-M (OSM)
Research Area:	Tumor Immunity
Clonality:	Polyclonal
Reactivity:	Mouse
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	OSM (Lys116-Arg205)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P53347 (UniProt , ExPASy)
Gene Symbol:	OSM
KEGG:	mmu:18413
String:	10090.ENSMUSP00000074708
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.