Apolipoprotein E (APOE) Antibody

Catalogue No.:abx101447



WB analysis of recombinant Human APOE protein.



WB analysis of Human liver lysate, using APOE Antibody (2 µg/ml).



WB analysis of Human serum, using APOE Antibody (2 µg/ml).



WB analysis of Mouse cerebrum lysate, using APOE Antibody (2 µg/ml).



WB analysis of Mouse lung lysate, using APOE Antibody (2 µg/ml).

Abbexa LTD, Cambridge, UK · Phone: +44 (0) 1223 755950 · Fax: +44 (0) 1223 755951
Abbexa LLC, Houston, TX USA · Phone: +1 832 327 7413
Abbexa BV, Leiden, NL
Website: www.abbexa.com · Email: info@abbexa.com



WB analysis of Mouse serum, using APOE Antibody (2 µg/ml).



IHC-P analysis of Human liver tissue, using APOE Antibody (20 µg/ml).



IHC-P analysis of Human kidney tissue, using APOE Antibody (20 µg/ml).



IHC-P analysis of Human cerebrum tissue, using APOE Antibody (20 µg/ml).



Polyclonal Antibody to Apolipoprotein E (APOE).

Abbexa LTD, Cambridge, UK · Phone: +44 (0) 1223 755950 · Fax: +44 (0) 1223 755951
Abbexa LLC, Houston, TX USA · Phone: +1 832 327 7413
Abbexa BV, Leiden, NL
Website: www.abbexa.com · Email: info@abbexa.com

Target:	Apolipoprotein E (APOE)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx065429 - Recombinant APOE (Lys19-Gln181) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P02649 (UniProt, ExPASy)
Gene Symbol:	APOE
GeneID:	348
OMIM:	104310
HGNC:	613
KEGG:	hsa:348
Ensembl:	ENSG00000130203
String:	9606.ENSP00000252486
Buffer:	PBS, pH 7.4., containing 0.02% NaN3, 50% glycerol.
Specificity:	Cross-reacts with Mouse and Pig APOE.
Concentration:	1 mg/ml

Abbexa LTD, Cambridge, UK · Phone: +44 (0) 1223 755950 · Fax: +44 (0) 1223 755951
Abbexa LLC, Houston, TX USA · Phone: +1 832 327 7413
Abbexa BV, Leiden, NL
Website: www.abbexa.com · Email: info@abbexa.com

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

Abbexa LTD, Cambridge, UK · Phone: +44 (0) 1223 755950 · Fax: +44 (0) 1223 755951
Abbexa LLC, Houston, TX USA · Phone: +1 832 327 7413
Abbexa BV, Leiden, NL
Website: www.abbexa.com · Email: info@abbexa.com