



Multifunctional Protein ADE2H1 (PAICS) Antibody

Catalogue No.:abx004943



Western blot analysis of various lysates using PAICS Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 30s.



Immunofluorescence analysis of U2OS cells using PAICS Antibody. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.

PAICS Antibody is a Rabbit Polyclonal antibody against PAICS. This gene encodes a bifunctional enzyme containing phosphoribosylaminoimidazole carboxylase activity in its N-terminal region and phosphoribosylaminoimidazole succinocarboxamide synthetase in its C-terminal region. It catalyzes steps 6 and 7 of purine biosynthesis. The gene is closely linked and divergently transcribed with a locus that encodes an enzyme in the same pathway, and transcription of the two genes is coordinately regulated. The human genome contains several pseudogenes of this gene. Multiple transcript variants encoding different isoforms have been found for this gene.

Target: Multifunctional Protein ADE2H1 (PAICS)

Clonality: Polyclonal

Reactivity: Human, Rat

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

Recommended dilutions: ELISA: 1 μg/ml, WB: 1/500 - 1/2000, IF/ICC: 1/10 - 1/100. Optimal dilutions/concentrations should

be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to PAICS. The exact sequence is proprietary.

Isotype: IgG

Datasheet

Version: 4.0.0 Revision date: 03 Sep 2025



Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P22234 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: PAICS

GeneID: <u>10606</u>

NCBI Accession: NP_001072992.1

KEGG: hsa:10606

String: <u>9606.ENSP00000382595</u>

Molecular Weight: Calculated MW: 47 kDa

Observed MW: 47 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.