



Serotonin (5-HT) Antibody


Catalogue No.: abx101820




IHC-P analysis with DAB staining of Rat small intestine tissue, using 5-HT antibody (20 µg/ml) and and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).




IHC-P analysis with DAB staining of Rat colon tissue, using 5-HT antibody (20 µg/ml) and and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis with DAB staining of Pig small intestine tissue, using 5-HT antibody (20 µg/ml) and and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).




IHC-P analysis with DAB staining of Pig colon tissue, using 5-HT antibody (20 µg/ml) and and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).




IHC-P analysis with DAB staining of Mouse colon tissue, using 5-HT antibody (10 µg/ml) and and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml). PBS was used instead of the primary antibody as a control.

Datasheet

Version: 6.0.0
Revision date: 27 Mar 2025



IHC-P analysis with DAB staining of Mouse small intestine tissue, using 5-HT antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml). PBS was used instead of the primary antibody as a control.



IHC-P analysis with DAB staining of Mouse stomach tissue, using 5-HT antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml). PBS was used instead of the primary antibody as a control.

Polyclonal Antibody to 5-Hydroxytryptamine.

Target:	Serotonin (5-HT)
Clonality:	Polyclonal
Reactivity:	General
Tested Applications:	ELISA, IHC
Host:	Rabbit
Recommended dilutions:	ELISA: 136 ng/ml, IHC: 5-20 µg/ml, IF/ICC (Predicted): 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx165617 - 5-HT conjugated to BSA
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Datasheet

Version: 6.0.0

Revision date: 27 Mar 2025



Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only