



MAPK/MAK/MRK Overlapping Kinase (MOK) Antibody


Catalogue No.:abx101916




Western blot analysis of Mouse Lung lysate, using Human RAGE Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of the recombinant protein.



IHC-P analysis of Human Liver cancer Tissue, with DAB staining.



IHC-P analysis of liver cancer tissue, with DAB staining.

Polyclonal Antibody to Renal Tumor Antigen (RAGE).

Target: MAPK/MAK/MRK Overlapping Kinase (MOK)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 2.0.0

Revision date: 28 Jun 2025



Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: RAGE (Leu173-Gly418)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q9UQ07 ([UniProt](#), [ExPASy](#))

String: [9606.ENSP00000355304](#)

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.