



Von Willebrand Factor (vWF) Antibody


Catalogue No.:abx102081




Western blot analysis of Rat Serum.



Western blot analysis of recombinant Rat vWF.



IHC-P analysis of Rat Pancreas Tissue, with DAB staining.



IHC-P analysis of Rat Pancreas Tissue, with DAB staining.

Polyclonal Antibody to vWF.

Target: Von Willebrand Factor (vWF)

Clonality: Polyclonal

Reactivity: Rat

Datasheet

Version: 2.0.0

Revision date: 06 Sep 2025



Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: vWF (Ser269-Pro367)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q62935 ([UniProt](#), [ExPASy](#))

String: [10116.ENSRNOP00000026643](#)

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.