



Muellerian-Inhibiting Factor (AMH) Antibody


Catalogue No.: abx102091




Western blot analysis of recombinant Human AMH.




Western blot analysis of (1) Rat Ovary Tissue and (2) Mouse Testis Tissue.




IHC-P analysis of Human Esophagus Cancer Tissue, with DAB staining, using Rabbit Anti-Human AMH Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).




IHC-P analysis of Human Liver Cancer Tissue, with DAB staining.



IHC-P analysis of Human Stomach Tissue, with DAB staining.



IHC-P analysis of Human Rectum Cancer Tissue, with DAB staining.



IHC-P analysis of Human Liver Tissue, with DAB staining.

Polyclonal Antibody to AMH (AMH).

Target:	Muellerian-Inhibiting Factor (AMH)
Research Area:	Endocrinology, Reproductive science, Hormone metabolism
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx065389 - Recombinant AMH (Ala453-Arg560) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Datasheet

Version: 4.0.0

Revision date: 10 Sep 2025



UniProt Primary AC: P03971 ([UniProt](#), [ExPASy](#))

Gene Symbol: AMH

GeneID: [268](#)

KEGG: hsa:268

String: [9606.ENSP00000221496](#)

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR **USE** IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only