



Tachykinin Receptor 2 (TACR2) Antibody


Catalogue No.: abx102297




Western blot analysis of Human Lung lysate, using Human TACR2 Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of Human Hela cell lysate, using Rabbit Anti-Human TACR2 Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of Pig Small intestine lysate, using Rabbit Anti-Human TACR2 Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of Porcine Heart lysate, using Rabbit Anti-Human TACR2 Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of the recombinant protein.

Datasheet

Version: 4.0.0
Revision date: 13 Sep 2025



IHC-P analysis of Hela Cells, with FITC staining.



IHC-P analysis of Human Pancreatic Cancer Tissue, with DAB staining.

Polyclonal Antibody to Tachykinin Receptor 2 (TACR2).

Target: Tachykinin Receptor 2 (TACR2)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: TACR2 (Val92-Leu290)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Datasheet

Version: 4.0.0

Revision date: 13 Sep 2025



Note:

THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only