



Prothrombin Fragment 1+2 (F1+2) Antibody


Catalogue No.: abx102478



Western blot analysis of recombinant Rat F1+2 Protein.



Western blot analysis of Rat Kidney lysate, using Rabbit Anti-Rat F1+2 Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of Rat Serum, using Rabbit Anti-Rat F1+2 Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of Rat Red Blood Cells lysate, using Rabbit Anti-Rat F1+2 Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of Rat Liver lysate, using Rabbit Anti-Rat F1+2 Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).

Datasheet

Version: 4.0.0
Revision date: 28 Sep 2025



Polyclonal Antibody to Prothrombin Fragment 1+2 (F1+2).

Target:	Prothrombin Fragment 1+2 (F1+2)
Clonality:	Polyclonal
Reactivity:	Rat
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-5 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx167765 - Recombinant F1+2 (Ala44-Arg323) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P18292 (UniProt , ExPASy)
String:	10116.ENSRNOP00000022233
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Specificity:	Cross-reacts with Mouse F1+2.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.