

Prothrombin Fragment 1+2 (F1+2) Antibody

Catalogue No.:abx102479



Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using Prothrombin Fragment 1+2 (F1+2) Antibody (20 μ g/ml) and HRP-conjugated Goat Anti-Rabbit antibody (abx400043, 2 μ g/ml), with DAB staining. PBS was used instead of the primary antibody for the negative control.



Immunohistochemistry analysis of paraffin-embedded Mouse lung tissue using Prothrombin Fragment 1+2 (F1+2) Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody (abx400043, 2 µg/ml), with DAB staining. PBS was used instead of the primary antibody for the negative control.





Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using Prothrombin Fragment 1+2 (F1+2) Antibody (20 μ g/ml) and HRP-conjugated Goat Anti-Rabbit antibody (<u>abx400043</u>, 2 μ g/ml), with DAB staining. PBS was used instead of the primary antibody for the negative control.



Immunofluorescence analysis of HepG2 cells using Prothrombin Fragment 1+2 (F1+2) Antibody (20 µg/ml) and AF488-conjugated Goat Anti-Rabbit antibody (2 µg/ml).

Prothrombin Fragment 1+2 (F1+2) Antibody is a rabbit Polyclonal Antibody to Prothrombin Fragment 1+2 (F1+2).

Target:	Prothrombin Fragment 1+2 (F1+2)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions	: WB: 0.01-2 μg/ml, IHC: 5-20 μg/ml, IF/ICC: 5-20 μg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx168232 - Recombinant Human F1+2 (Ala44-Arg327) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

abbexa 🔶

Stability:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.
UniProt Primary AC:	P00734 (<u>UniProt</u> , <u>ExPASy</u>)
KEGG:	hsa:2147
String:	9606.ENSP00000308541
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Specificity:	Cross-reacts with Mouse, Rat and Pig F1+2.
Concentration:	0.5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.