



Indoleamine-2,3-Dioxygenase (IDO) Antibody


Catalogue No.: abx102654



Western blot analysis of Rat Placenta Tissue.



Western blot analysis of recombinant Human IDO.



IHC-P analysis of Human Tissue, with DAB staining.

Polyclonal Antibody to Indoleamine-2, 3-Dioxygenase (IDO).

Target: Indoleamine-2,3-Dioxygenase (IDO)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1-5 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 2.0.0

Revision date: 27 Jun 2025



Immunogen:	IDO (Ala2-Gly403)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P14902 (UniProt , ExPASy)
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only