



Histone Cluster 1, H2ag (HIST1H2AG) Antibody


Catalogue No.: abx102875



Western blot analysis of recombinant Human HIST1H2AG.



IHC-P analysis of Human Stomach Tissue, with DAB staining.



IHC-P analysis of Human Prostate Gland Tissue, with DAB staining.

Polyclonal Antibody to Histone Cluster 1, H2ag (HIST1H2AG).

Target: Histone Cluster 1, H2ag (HIST1H2AG)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 2.0.0

Revision date: 11 Oct 2025



Immunogen:	HIST1H2AG (Met1-Lys130)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only