



Tumor Necrosis Factor Ligand Superfamily Member 6 (FASLG) Antibody

Catalogue No.: abx103238



Western blot analysis of the recombinant protein.



IHC-P analysis of Human Tissue, with DAB staining.

Polyclonal Antibody to Tumor Necrosis Factor Ligand Superfamily Member 6 (FASLG).

Target: Tumor Necrosis Factor Ligand Superfamily Member 6 (FASLG)

Research Area: Apoptosis, Tumour Immunity, Infection Immunity

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: FASLG (Gln103-Leu281)

Form: Liquid

Datasheet

Version: 3.0.0

Revision date: 04 Oct 2025



Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P48023 (UniProt , ExPASy)
Gene Symbol:	FASLG
KEGG:	hsa:356
String:	9606.ENSP00000356694
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only