



Tumor Necrosis Factor Ligand Superfamily Member 6 (FASLG) Antibody


Catalogue No.:abx103240



Western blot analysis of recombinant Rat FASLG.




IHC-P analysis of Rat Kidney Tissue, with DAB staining.



IHC-P analysis of Rat Pancreas Tissue, with DAB staining.



IHC-P analysis of Rat Uterus Tissue, with DAB staining.



IHC-P analysis of Rat Pancreas Tissue, with DAB staining.

Datasheet

Version: 4.0.0 Revision date: 12 Sep 2025



Polyclonal Antibody to Tumor Necrosis Factor Ligand Superfamily Member 6 (FASLG).

Target: Tumor Necrosis Factor Ligand Superfamily Member 6 (FASLG)

Research Area: Apoptosis, Tumour Immunity, Infection Immunity

Clonality: Polyclonal

Reactivity: Rat

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be

determined by the end user.

Conjugation: Unconjugated

Immunogen: FASLG (Gln100-Leu278)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P36940 (UniProt, ExPASy)

Gene Symbol: FASLG

GeneID: <u>25385</u>

KEGG: rno:25385

String: <u>10116.ENSRNOP00000003998</u>

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.