



Heparan Sulfate Proteoglycan 2 (HSPG2) Antibody


Catalogue No.: abx103270




Western blot analysis of recombinant Human HSPG2 protein.




Western blot analysis of Pig Heart lysate, using Rabbit Anti-Human HSPG2 Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of Mouse Lung lysate, using Rabbit Anti-Human HSPG2 Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




IHC-P analysis of Human Liver Cancer Tissue, with DAB staining, using Rabbit Anti-Human HSPG2 Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).




IHC-P analysis of Pig Stomach Tissue, with DAB staining, using Rabbit Anti-Human HSPG2 Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Pig Spleen Tissue, with DAB staining, using Rabbit Anti-Human HSPG2 Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Pig Kidney Tissue, with DAB staining, using Rabbit Anti-Human HSPG2 Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Pig Cardiac Muscle Tissue, with DAB staining, using Rabbit Anti-Human HSPG2 Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Polyclonal Antibody to Heparan Sulfate Proteoglycan 2 (HSPG2).

Target: Heparan Sulfate Proteoglycan 2 (HSPG2)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-5 µg/ml, IHC: 5-50 µg/ml, IF/ICC: 5-50 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx067054](#) - Recombinant HSPG2 (Leu4149-Ser4391) expressed in E. coli

Datasheet

Version: 3.0.0
Revision date: 08 Oct 2025



Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P98160 (UniProt , ExPASy)
KEGG:	hsa:3339
String:	9606.ENSP00000363827
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Specificity:	Cross-reacts with Mouse, Rat and Pig HSPG2.
Concentration:	0.67 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only