



Activating Transcription Factor 7 (ATF7) Antibody


Catalogue No.:abx103396




Western blot analysis of (1) Human SGC7901 Cells, (2) Human HeLa cells and (3) Rat Stomach Tissue).




Western blot analysis of recombinant Mouse ATF7.



IHC-P analysis of Mouse Ovary Tissue, with DAB staining.




IHC-P analysis of Mouse Kidney Tissue, with DAB staining.



IHC-P analysis of Mouse Spleen Tissue, with DAB staining.

Datasheet

Version: 3.0.0
Revision date: 07 Oct 2025



IHC-P analysis of Mouse Intestine Tissue, with DAB staining.

Polyclonal Antibody to Activating Transcription Factor 7 (ATF7).

Target:	Activating Transcription Factor 7 (ATF7)
Clonality:	Polyclonal
Reactivity:	Mouse
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	ATF7 (Gln50-Ser270)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.