



C-X3-C Motif Chemokine Ligand 1 (CX3CL1) Antibody


Catalogue No.: abx103407




Western blot analysis of (1) Mouse Brain Tissue and (2) Mouse Heart Tissue.




Western blot analysis of recombinant Human CX3CL1.




IHC-P analysis of Human Prostate Gland Tissue, with DAB staining.




IHC-P analysis of Human Breast Cancer Tissue, with DAB staining.




IHC-P analysis of Human Stomach Cancer Tissue, with DAB staining.



IHC-P analysis of Human Kidney Tissue, with HE staining.



IHC-P analysis of Human Testis Tissue, with HE staining.



IHC-P analysis of Human Liver Tissue, with DAB staining.

Polyclonal Antibody to CX3CL1.

Target: C-X3-C Motif Chemokine Ligand 1 (CX3CL1)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-5 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: CX3CL1 (Gln25-Pro332)

Datasheet

Version: 2.0.0

Revision date: 03 Apr 2025



Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P78423 (UniProt , ExPASy)
KEGG:	hsa:6376
String:	9606.ENSP00000006053
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only